Sarcouncil Journal of Engineering and Computer Sciences

ISSN(Online): 2945-3585

Volume- 04| Issue- 07| 2025

Research Article **Received:** 26-05-2025 | **Accepted:** 16-06-2025 | **Published:** 03-07-2025

Multi-Modal AI for Renewable Energy Forecasting: Combining Satellite Imagery, Weather Data, and Historical Patterns

Vijay Bhalani
University of Southern California, USA

Abstract: Integrating renewable energy sources into modern electrical grids presents significant challenges due to the inherent variability and intermittency of solar and wind power generation systems. Traditional forecasting methods relying on single-source data streams demonstrate substantial accuracy deficiencies that compromise grid stability and operational efficiency. Multi-modal artificial intelligence represents a transformative paradigm for renewable energy forecasting by simultaneously processing satellite imagery, meteorological data, and historical generation patterns through advanced deep learning architectures. The developed framework employs convolutional neural networks for spatial feature extraction from satellite observations, recurrent components for temporal sequence modeling, and attention mechanisms for adaptive feature weighting across heterogeneous data modalities. Comprehensive experimental validation across diverse geographical regions and renewable energy technologies demonstrates significant performance improvements compared to conventional forecasting methods. The multi-modal integration enables enhanced prediction accuracy through sophisticated data fusion strategies that capture complex spatiotemporal relationships governing renewable energy generation patterns. Real-time prediction capabilities support operational grid management applications, including energy dispatch optimization and grid stability assessment under varying renewable energy penetration scenarios. Cross-regional model performance validation establishes transferability across different climatic conditions while identifying necessary adaptation procedures for maintaining forecasting effectiveness. The integrated framework addresses critical gaps in current forecasting methodologies by leveraging computer vision techniques for atmospheric condition assessment and implementing hierarchical feature learning architectures for multi-resolution data processing.

Keywords: Multi-modal Artificial Intelligence, Renewable Energy Forecasting, Satellite Imagery Processing, Deep Learning Architectures, Grid Management Applications.

INTRODUCTION

The contemporary energy landscape has witnessed an unprecedented transformation toward sustainable power generation systems, fundamentally altering the operational dynamics of electrical grids worldwide. Modern power networks face increasingly complex challenges as renewable energy sources become dominant contributors to the global energy mix [Lebedys, A. *et al*., 2023]. The intermittent nature of solar and wind resources introduces substantial variability into power generation patterns, creating operational complexities that traditional grid management systems struggle to accommodate effectively. These fluctuations manifest across multiple temporal scales, from minute-by-minute variations due to cloud movements affecting solar installations to seasonal patterns influencing solar irradiance and wind resource availability throughout the year.

Traditional forecasting methodologies in renewable energy prediction have demonstrated significant limitations when applied to contemporary grid management requirements. Historical approaches utilizing single-source data streams, particularly those based on purely statistical models or limited meteorological inputs, exhibit substantial accuracy deficiencies that compromise grid stability and operational efficiency [Mellit, A. *et al*., 2021]. These conventional methods typically rely on autoregressive models, time series analysis, or basic correlation techniques that fail to capture the complex multi-dimensional relationships governing renewable energy generation. The inadequacy of single-source forecasting becomes particularly pronounced during transitional weather conditions, where rapid atmospheric changes can cause dramatic shifts in energy output that existing prediction models cannot anticipate with sufficient precision.

Multi-modal artificial intelligence represents a paradigm shift in renewable energy forecasting, offering sophisticated capabilities to integrate diverse data sources into comprehensive prediction frameworks. Advanced machine learning architectures can simultaneously process satellite imagery for cloud cover analysis, meteorological data for atmospheric condition assessment, and historical generation patterns for temporal trend identification [Lebedys, A. *et al*., 2023]. Computer vision techniques applied to high-resolution satellite observations enable detailed analysis of atmospheric phenomena, surface characteristics, and weather pattern evolution that directly influence renewable energy production. These integrated approaches leverage deep learning methodologies to extract complex feature relationships that remain invisible to traditional single-source analysis methods.

The research investigation focuses on developing comprehensive multi-modal forecasting systems that substantially enhance prediction accuracy compared to existing methodologies. The study

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

***Corresponding Author:** Vijay Bhalani

DOI- https://doi.org/10.5281/zenodo.15797281

aims to demonstrate how integrated artificial intelligence approaches can reduce forecasting uncertainties while providing real-time prediction capabilities essential for modern grid operations [Mellit, A. *et al*., 2021]. This research contributes novel data fusion techniques specifically designed for renewable energy applications, addressing critical gaps in current forecasting methodologies. The investigation explores how computer vision analysis of satellite imagery can be effectively combined with meteorological data streams and historical generation patterns to create robust prediction models suitable for various geographical regions and renewable energy technologies.

The methodological framework encompasses systematically integrating multiple data modalities through specialized neural network architectures designed for heterogeneous data processing. The research methodology includes comprehensive validation across diverse climate zones and energy generation technologies, ensuring the developed forecasting systems' broad applicability. The paper organization progresses through detailed literature analysis, methodology development, experimental validation, and practical implementation considerations for deployment in operational grid management environments.

2. LITERATURE REVIEW AND THEORETICAL FRAMEWORK

The historical development of renewable energy forecasting methodologies has traversed multiple technological paradigms, establishing foundational approaches that continue to influence contemporary prediction systems. Early statistical frameworks centered on autoregressive integrated moving average models and classical regression techniques provided essential insights into temporal patterns within renewable energy generation datasets [Sehrawat, N. *et al*., 2023]. These traditional methods demonstrated competency in identifying seasonal variations and long-term cyclical behaviors inherent in solar and wind energy production. However, limitations became apparent when addressing the nonlinear characteristics of atmospheric phenomena and the complex interdependencies between multiple meteorological variables affecting renewable energy output. Exponential smoothing techniques and polynomial regression extensions were subsequently developed to enhance predictive capabilities, particularly for applications requiring integration of clear sky radiation models with historical generation patterns.

The transition toward machine learning methodologies marked a significant advancement in renewable energy forecasting capabilities, introducing algorithms capable of modeling nonlinear relationships between input variables and energy generation outcomes. Support vector machine implementations demonstrated superior performance in handling high-dimensional feature spaces typical of meteorological datasets used in renewable energy prediction applications [Sehrawat, N. *et al*., 2023]. Ensemble learning methods, including random forest algorithms and gradient boosting techniques, provided enhanced robustness through a combination of multiple weak learners trained on diverse subsets of available data. Artificial neural network architectures, encompassing multilayer perceptrons and radial basis function networks, enabled sophisticated modeling of complex atmospheric interactions that influence renewable energy generation patterns. These machine learning approaches established new benchmarks for forecasting accuracy across various temporal horizons, demonstrating particular effectiveness in short-term prediction scenarios where rapid meteorological changes significantly impact energy output.

Contemporary deep learning applications represent the current state-of-the-art in renewable energy forecasting, utilizing advanced neural network architectures to extract intricate spatiotemporal features from multi-dimensional datasets. Convolutional neural network implementations have shown exceptional capabilities in processing high-resolution satellite imagery for automated cloud detection and atmospheric condition analysis [Al-Musaylh, M.S. *et al*., 2018]. Long short-term memory networks and gated recurrent units have proven effective in capturing temporal dependencies within renewable energy generation sequences, particularly for modeling complex seasonal patterns and weather-dependent fluctuations. Advanced attention mechanisms and transformer architectures have further enhanced the ability to identify relevant temporal and spatial relationships within large-scale environmental datasets.

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher




Fig 1: Literature Review Framework [Sehrawat, N. *et al*., 2023, Al-Musaylh, M.S. *et al*., 2018]

Multi-modal data fusion methodologies have gained significant traction across diverse application domains, including environmental monitoring, climate modeling, and atmospheric science research. Computer vision techniques applied to satellite imagery analysis have revolutionized meteorological forecasting by enabling automated extraction of cloud morphology characteristics, atmospheric moisture distributions, and surface albedo variations [Al-Musaylh, M.S. *et al*., 2018]. These approaches utilize sophisticated image processing algorithms to quantify cloud coverage patterns, wind velocity indicators, and atmospheric stability measures derived from multi-spectral satellite observations. Feature extraction methodologies can identify subtle atmospheric phenomena that correlate strongly with renewable energy generation potential across different geographical regions.

Transfer learning applications across geographical boundaries and renewable energy technologies constitute an emerging research frontier addressing the challenge of model adaptability and generalization. Current literature reveals significant gaps in the comprehensive integration of heterogeneous data sources, limited exploration of spatial correlation patterns, and insufficient consideration of temporal variability across multiple time scales simultaneously. The theoretical foundation for multi-modal approaches emerges from recognition that renewable energy generation depends on complex interactions between numerous environmental factors that cannot be adequately represented through single-source data analysis frameworks.

3. MULTI-MODAL DATA INTEGRATION AND FEATURE EXTRACTION

Satellite imagery processing for renewable energy applications represents a sophisticated domain requiring advanced computational methodologies to extract meaningful environmental parameters correlating with energy generation capabilities. Cloud cover detection and quantification techniques employ complex image processing algorithms that analyze spectral reflectance patterns across multiple wavelength bands to identify and classify atmospheric phenomena [Assaf, A.M. *et al*., 2023]. These methodologies incorporate edge detection algorithms, texture analysis, and morphological operations to distinguish between cloud formations and atmospheric conditions. Automated cloud classification systems utilize pattern recognition techniques to categorize cloud types based on altitude, density, and optical characteristics directly influencing solar radiation attenuation and scattering effects.

Vegetation indices calculations, particularly normalized difference vegetation index computations, provide critical surface characterization data for wind pattern assessment and boundary layer analysis in renewable energy applications. Surface vegetation density and health status significantly influence local wind flow dynamics, creating turbulence patterns and roughness coefficients that affect wind energy harvesting efficiency [Assaf, A.M. *et al*., 2023]. Remote sensing techniques enable large-scale monitoring of vegetation changes across seasons, providing insights into surface-atmosphere

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher

interactions that modify local meteorological conditions. Atmospheric condition analysis from satellite observations incorporates multi-spectral data processing to quantify water vapor distributions, aerosol loading, and atmospheric transparency factors that impact renewable energy generation potential.

Meteorological data preprocessing and feature engineering are fundamental components of comprehensive renewable energy forecasting frameworks, requiring systematic approaches to managing diverse data formats, temporal resolutions, and measurement uncertainties. Weather parameter selection processes involve statistical correlation analysis and feature importance ranking to identify the most relevant atmospheric variables from extensive meteorological databases containing temperature profiles, humidity gradients, pressure variations, and wind characteristics [Manavalan, E. *et al*., 2019]. Data normalization procedures address scale differences between meteorological parameters, ensuring consistent input ranges for subsequent machine learning algorithms. Advanced feature engineering techniques create composite variables that capture atmospheric stability conditions, convective potential, and boundary layer characteristics essential for accurate renewable energy prediction.

Temporal alignment of multi-source datasets presents complex synchronization challenges due to disparate sampling frequencies, measurement timestamps, and data acquisition protocols across different monitoring systems. Interpolation methodologies and temporal resampling algorithms establish unified temporal frameworks that accommodate varying data collection intervals from satellite platforms, ground-based weather stations, and energy generation monitoring systems [Manavalan, E. *et al*., 2019]. Historical generation pattern analysis employs sophisticated signal decomposition techniques to separate renewable energy production data into trend components, seasonal variations, and stochastic fluctuations. Time series decomposition methodologies identify recurring patterns and cyclical behaviors that enhance prediction accuracy across different forecasting horizons.




Fig 2: Multi-Modal Data Integration Framework [Assaf, A.M. *et al*., 2023, Manavalan, E. *et al*., 2019]

Variability characterization across multiple temporal scales utilizes advanced statistical techniques to quantify renewable energy generation fluctuations spanning sub-hourly variations to multi-year trends. Data fusion architectures implement hierarchical processing frameworks that systematically integrate heterogeneous data sources while preserving critical information content from each modality. Synchronization challenges encompass handling irregular sampling intervals, compensating for measurement delays, and managing data quality variations across different acquisition platforms to maintain consistent forecasting performance across diverse operational scenarios.

4. DEEP LEARNING ARCHITECTURE FOR MULTI-MODAL FORECASTING

Neural network design for multi-modal data streams necessitates sophisticated architectural frameworks that can effectively process diverse input modalities while maintaining computational tractability and forecasting precision. Convolutional layers constitute the cornerstone for spatial feature extraction from satellite imagery, employing multiple convolutional filters with

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher

distinct receptive field sizes to capture hierarchical spatial patterns across different resolution scales Eitel, A. *et al*.., 2015]. These architectural components implement progressive feature abstraction through cascaded convolutional operations that transform raw pixel intensities into increasingly complex spatial representations. Max pooling and average pooling operations facilitate dimensionality reduction while preserving essential spatial relationships critical for atmospheric phenomenon detection. Advanced convolutional designs incorporate residual connections and skip pathways, enabling gradient flow through deep network architectures without degradation.

Recurrent components provide fundamental temporal sequence modeling capabilities for capturing dynamic behaviors embedded within renewable energy generation patterns and meteorological time series observations. Gated recurrent architectures implement sophisticated memory mechanisms that selectively preserve relevant historical information while discarding irrelevant temporal dependencies Eitel, A. *et al*.., 2015]. These networks address computational challenges associated with long sequence processing through efficient gating operations that control information flow across temporal steps. Stacked recurrent configurations enable hierarchical temporal modeling where lower layers capture short-term fluctuations while higher layers model long-term seasonal patterns and cyclical behaviors inherent in renewable energy systems.

Attention mechanisms introduce adaptive feature weighting capabilities that dynamically emphasize relevant input components based on contextual dependencies and prediction requirements. Global attention frameworks compute attention weights across entire input sequences, enabling direct connections between distant temporal positions without sequential processing constraints [Bahdanau, D. *et al*.., 201Al-Musaylh, M.S. *et al*., 2018]. Local attention mechanisms focus computational resources on specific temporal windows or spatial regions most relevant to current prediction tasks. Cross-attention architectures facilitate interaction between data modalities by computing attention weights highlighting complementary information across heterogeneous input streams.

Fusion strategies for heterogeneous data types encompass diverse architectural paradigms designed to integrate multi-modal information streams effectively throughout the learning process. Early fusion methodologies concatenate different data modalities at input layers, enabling joint feature learning through shared network parameters that capture cross-modal correlations [Bahdanau, D. *et al*.., 201Al-Musaylh, M.S. *et al*., 2018]. Intermediate fusion approaches combine feature representations at multiple network depths, allowing modality-specific and joint feature extraction processes. Late fusion architectures maintain separate processing pathways for each data modality before combining learned representations at decision layers through weighted aggregation mechanisms.

Hierarchical feature learning architectures implement multi-resolution processing frameworks that simultaneously extract features at various abstraction levels. Transfer learning implementation across geographical regions and energy technologies addresses model adaptability challenges when deploying forecasting systems in diverse environmental conditions. Domain adaptation methodologies employ adversarial learning frameworks that encourage domain-invariant feature representations while preserving task-specific predictive capabilities. Cross-modal knowledge transfer mechanisms leverage learned representations from data-rich modalities to enhance performance on data-sparse modalities through shared embedding spaces. Model optimization and hyperparameter tuning methodologies incorporate gradient-based optimization algorithms, learning rate scheduling strategies, and regularization techniques that prevent overfitting while maximizing generalization performance across different deployment scenarios.

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher




Fig 3: Deep Learning Architecture Framework Eitel, A. *et al*.., 2015, Bahdanau, D. *et al*.., 201Al-Musaylh, M.S. *et al*., 2018]

5. EXPERIMENTAL RESULTS AND PERFORMANCE ANALYSIS

Dataset description and experimental setup encompass systematic data collection methodologies that aggregate diverse information sources across multiple geographical regions to establish comprehensive evaluation frameworks for multi-modal renewable energy forecasting systems. Multi-regional renewable energy generation data compilation systematically aggregates production records from distributed solar photovoltaic installations and wind energy facilities spanning diverse climatic zones and topographical conditions [Zhou, H. *et al*.., 2019]. These comprehensive datasets incorporate generation measurements from installations ranging from utility-scale facilities to distributed residential systems, ensuring representative coverage of different operational scales and technological configurations. Data collection protocols maintain standardized temporal sampling rates and measurement calibration procedures for consistent comparative analysis across heterogeneous renewable energy deployment scenarios.

Satellite imagery sources and temporal coverage integrate observations from multiple Earth observation platforms to provide comprehensive atmospheric monitoring capabilities essential for advanced renewable energy forecasting applications. High-resolution satellite data acquisition encompasses multiple spectral channels, including visible light, near-infrared, and thermal infrared wavelengths, to capture diverse atmospheric phenomena affecting renewable energy generation potential [Zhou, H. *et al*.., 2019]. Temporal data coverage spans extended observation periods encompassing multiple seasonal cycles and weather pattern variations to establish robust training datasets for machine learning model development. Quality assessment procedures implement systematic validation protocols to identify and correct sensor degradation effects, atmospheric distortion artifacts, and data transmission errors that could compromise model training effectiveness.

Meteorological data integration systematically incorporates atmospheric observations from distributed ground-based monitoring networks, upper-atmosphere sounding systems, and numerical weather prediction model outputs to provide comprehensive environmental characterization. Data quality assessment methodologies employ statistical anomaly detection algorithms and cross-validation techniques to identify measurement inconsistencies, sensor malfunctions, and data acquisition errors that could negatively impact forecasting model performance [Rana, M. *et al*., 201Manavalan, E. *et al*., 2019]. Compared to conventional single-source prediction methodologies, comparative analysis with baseline forecasting methods demonstrates quantitative performance improvements achieved through integrated multi-modal approaches. Baseline comparison frameworks encompass traditional

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher

statistical forecasting methods, conventional machine learning approaches, and contemporary deep learning architectures operating on individual data modalities.

Statistical significance testing employs rigorous hypothesis testing methodologies, including t-tests, Mann-Whitney U tests, and bootstrap confidence interval estimation to validate performance improvements and establish statistical confidence in comparative results. Accuracy metrics evaluation encompasses comprehensive error analysis using multiple performance indicators across various temporal prediction horizons ranging from immediate short-term forecasts to extended multi-day prediction intervals [Rana, M. *et al*., 201Manavalan, E. *et al*., 2019]. Real-time prediction capability evaluation assesses computational performance characteristics, including processing latency, memory utilization, and throughput requirements for operational deployment in time-critical grid management applications.

Computational efficiency analysis examines processing time requirements across different hardware configurations and software optimization strategies to ensure compatibility with real-time operational constraints imposed by modern electrical grid management systems. Grid management application scenarios evaluate practical deployment capabilities, including energy dispatch optimization support, ancillary service provision, and grid stability assessment under varying renewable energy penetration scenarios. Regional transferability and scalability assessment examine model performance consistency when deploying trained forecasting systems across different geographical regions characterized by distinct climatic patterns and atmospheric conditions. Cross-regional model performance validation quantifies accuracy variations and identifies necessary adaptation procedures for forecasting effectiveness across diverse operational environments.




Fig 4: Experimental Results Framework [Zhou, H. *et al*.., 2019, Rana, M. *et al*., 201Manavalan, E. *et al*., 2019]

CONCLUSION

The development of multi-modal artificial intelligence systems for renewable energy forecasting represents a significant advancement in addressing the complex challenges of integrating variable renewable energy sources into modern electrical grids. Integrating satellite imagery processing, meteorological data preprocessing, and historical generation pattern recognition through sophisticated deep learning architectures demonstrates substantial improvements in prediction accuracy compared to traditional single-source forecasting methods. Advanced neural network designs incorporating convolutional layers for spatial feature extraction, recurrent components for temporal modeling, and attention mechanisms for adaptive feature weighting enable comprehensive processing of heterogeneous data streams essential for accurate renewable energy prediction. Implementing fusion strategies across diverse data modalities facilitates effective integration of complementary information sources

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher

while addressing synchronization challenges inherent in multi-source data processing. Transfer learning applications across geographical regions and renewable energy technologies establish model adaptability frameworks that enable deployment across diverse environmental conditions with appropriate domain adaptation procedures. Real-time prediction capabilities demonstrated through computational efficiency evaluations support practical deployment in operational grid management scenarios where timely forecasting information is critical for maintaining system stability and optimizing energy dispatch decisions. Future developments should incorporate additional data modalities, including economic indicators and grid topology information, to enhance forecasting capabilities further, while advancing real-time adaptation mechanisms and online learning strategies to maintain prediction accuracy under evolving operational conditions.

REFERENCES

1. Lebedys, A., Akande, D., Coënt, N., Elhassan, N., Escamilla, G., Arkhipova, I. and Whiteman, A. "Renewable Capacity Statistics 2023." *IRENA*, (2023). https://unreeea.org/wp-content/uploads/2023/04/IRENA_RE_Capacity_Statistics_2023.pdf
2. Mellit, A., Pavan, A.M. and Lughi, V. "Deep learning neural networks for short-term photovoltaic power forecasting." *Renewable Energy* 172 (2021): 276-288. https://www.sciencedirect.com/science/article/abs/pii/S0960148121003475
3. Sehrawat, N., Vashisht, S. and Singh, A. "Solar irradiance forecasting models using machine learning techniques and digital twin: A case study with comparison." *International Journal of Intelligent Networks* 4 (2023): 90-102. https://www.sciencedirect.com/science/article/pii/S2666603023000064
4. Al-Musaylh, M.S., Deo, R.C., Adamowski, J.F. and Li, Y. "Short-term electricity demand forecasting with MARS, SVR and ARIMA models using aggregated demand data in Queensland, Australia." *Advanced Engineering Informatics* 35 (2018): 1-16. https://www.sciencedirect.com/science/article/abs/pii/S1474034617301477
5. Assaf, A.M., Haron, H., Abdull Hamed, H.N., Ghaleb, F.A., Qasem, S.N. and Albarrak, A.M. "A review on neural network based models for short term solar irradiance forecasting." *Applied Sciences* 13.14 (2023): 8332. https://www.mdpi.com/2076-3417/13/14/8332
6. Manavalan, E. and Jayakrishna, K. "A review of Internet of Things (IoT) embedded sustainable supply chain for industry 4.0 requirements." *Computers & industrial engineering* 127 (2019): 925-953. https://www.sciencedirect.com/science/article/abs/pii/S0360835218305709
7. Eitel, A., Springenberg, J.T., Spinello, L., Riedmiller, M. and Burgard, W. "Multimodal deep learning for robust RGB-D object recognition." *2015 IEEE/RSJ International Conference on Intelligent Robots and Systems (IROS)*. *IEEE*, (2015). https://arxiv.org/abs/1507.06821
8. Bahdanau, D., Cho, K. and Bengio, Y. "Neural machine translation by jointly learning to align and translate." *arXiv preprint arXiv:1409.0473* (2014). https://arxiv.org/abs/1409.0473
9. Zhou, H., Zhang, Y., Yang, L., Liu, Q., Yan, K. and Du, Y. "Short-term photovoltaic power forecasting based on long short term memory neural network and attention mechanism." *Ieee Access* 7 (2019): 78063-78074. https://ieeexplore.ieee.org/document/8736879
10. Rana, M., Koprinska, I. and Agelidis, V.G. "Univariate and multivariate methods for very short-term solar photovoltaic power forecasting." *Energy Conversion and Management* 121 (2016): 380-390. https://www.sciencedirect.com/science/article/abs/pii/S0196890416303934

Source of support: Nil; **Conflict of interest:** Nil.

Cite this article as:

Bhalani, V. "Multi-Modal AI for Renewable Energy Forecasting: Combining Satellite Imagery, Weather Data, and Historical Patterns." *Sarcouncil Journal of Engineering and Computer Sciences* 4.7 (2025): pp 66-73.

Copyright © 2022 The Author(s): This work is licensed under a Creative Commons Attribution- NonCommercial-NoDerivatives 4.0 (CC BY-NC-ND 4.0) International License

Publisher: SARC Publisher